

Fifteen most common *Salmonella* serovars from human sources in Japan, 2005-2009

(Infectious Agents Surveillance Report: Data based on the reports received before November 4, 2010 from public health institutes)

2009		2008		2007		2006		2005	
Enteritidis	232	Enteritidis	346	Enteritidis	585	Enteritidis	382	Enteritidis	725
Infantis	90	Infantis	107	Typhimurium	98	Typhimurium	73	Infantis	79
Thompson	69	Typhimurium	85	Thompson	85	Infantis	67	Typhimurium	63
Saintpaul	68	Saintpaul	70	Montevideo	83	Saintpaul	65	Thompson	61
Typhimurium	53	Braenderup	65	Saintpaul	75	Thompson	43	Montevideo	50
Bareilly	32	Thompson	60	Infantis	72	Newport	33	Litchfield	35
Schwarzengrund	28	Montevideo	50	Braenderup	52	Agona	31	Saintpaul	34
Montevideo	27	Typhi	28	Litchfield	29	Typhi	27	Agona	23
I 4:i:-	18	Stanley	23	Typhi	22	Litchfield	25	Derby	20
Litchfield	15	Litchfield	19	Newport	22	Montevideo	20	Braenderup	20
Manhattan	14	Schwarzengrund	18	Schwarzengrund	21	Virchow	20	Bareilly	16
Poona	12	Newport	16	Agona	20	Hadar	20	Virchow	16
Paratyphi B	11	Paratyphi A	12	Stanley	17	Stanley	16	Hadar	15
Newport	11	Agona	12	I 4:i:-	17	Paratyphi B	16	Schwarzengrund	12
Nagoya	11	Nagoya	12	Bareilly	17	Corvallis	13	Oranienburg	12
Others	202	Others	215	Hadar	17	Others	311	Anatum	12
Total	893	Total	1138	Others	300	Total	1162	Others	353
				Total	1532			Total	1546



Infectious Agents Surveillance Report