

【Table 1.】 Number of reports and percentage* about each symptom and medical intervention by age group (as of 24-Feb) *Excluded incomplete or unknown status cases from denominator

range	Fever (n=112)	Cough (n=112)	Pneumonia (n=66)	Sore throat (n=68)	General malaise (n=61)	Nasal discharge or congestion (n=64)	Headache (n=62)	Diarrhea (n=64)	Nausea and/or Vomiting (n=60)	Joint or Muscle pain (n=58)	Acute Respiratory Distress Syndrome (ARDS) (n=41)	Conjunctival congestion (n=53)	ICU Admission (n=52)	Invasive ventilation (n=50)
10-19	0 (0%)	0 (0%)	-	-	-	-	-	-	-	-	-	-	0 (0%)	0 (0%)
20-29	5 (71%)	6 (86%)	2 (40%)	1 (20%)	0 (0%)	3 (50%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
30-39	8 (100%)	5 (63%)	3 (75%)	2 (50%)	1 (25%)	0 (0%)	0 (0%)	1 (25%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
40-49	9 (100%)	5 (56%)	4 (80%)	2 (33%)	2 (40%)	0 (0%)	2 (40%)	1 (20%)	0 (0%)	1 (25%)	0 (0%)	0 (0%)	0 (0%)	1 (20%)
50-59	15 (75%)	15 (75%)	7 (64%)	5 (45%)	3 (30%)	3 (33%)	4 (33%)	3 (27%)	1 (10%)	1 (10%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
60-69	11 (85%)	6 (46%)	7 (78%)	4 (44%)	4 (50%)	2 (25%)	4 (50%)	1 (13%)	1 (11%)	0 (0%)	1 (14%)	0 (0%)	2 (25%)	2 (25%)
70-79	25 (76%)	18 (55%)	14 (67%)	5 (28%)	6 (38%)	5 (26%)	5 (31%)	3 (17%)	3 (18%)	2 (13%)	1 (9%)	0 (0%)	4 (24%)	5 (31%)
80-89	8 (40%)	14 (70%)	6 (55%)	4 (27%)	4 (29%)	4 (27%)	2 (14%)	2 (14%)	0 (0%)	0 (0%)	1 (13%)	0 (0%)	2 (29%)	3 (50%)